

FACTORIZATION OF THE NONLINEAR SCHRÖDINGER EQUATION AND APPLICATIONS

SWANHILD BERNSTEIN

ABSTRACT. We consider factorizations of the stationary and non-stationary Schrödinger equation in $\mathbb{R}^n$ which are based on appropriate Dirac operators. These factorizations lead to a Miura transform which is an analogue of the classical one-dimensional Miura transform but also closely related to the Riccati equation. In fact, the Miura transform is a nonlinear Dirac equation. We give an iterative procedure which is based on fix-point principles to solve this nonlinear Dirac equation. The relationship to nonlinear Schrödinger equations like the Gross-Pitaevskii equation are highlighted.

Keywords Clifford analysis, Schrödinger equation, nonlinear Dirac equation

Mathematical Subject Classification. Primary 30G35, Secondary 35J10, 35F30, 35Q55.

1. INTRODUCTION

Also today *nonlinear* equations and operators are a challenge. But it is also well-known that transformations do a good job in understanding and solving nonlinear equations. One famous transform is the Miura transform ([27], 1968) which in one spatial dimension represents a connection between Korteweg-de Vries equations. The Miura transform can also be considered as a special case of the Riccati equation and can be used to factorize the (stationary) Schrödinger equation. In higher dimensions Clifford algebras and analysis give the possibility to generalize the factorizations into higher spatial dimensions and even to space-time domains $(t, x) \in \mathbb{R}_+ \times G \subset \mathbb{R}_+ \times \mathbb{R}^n$.

Factorization of second order differential operators have been studied several times. Factorizations of the Helmholtz operator in case of a real wave number had been studied by K. Gürlebeck [11] and K. Gürlebeck/W. Sprößig in their books [12], [13]. Z. Xu [33], [34], F. Brackx and N. van Acker [6] and together with R. Delanghe and F. Sommen [7] considered generalized Dirac operators of type $D + k$, with the Dirac operator D and k is a complex number. E. Obolashvili [28], [29] and later on Huang Liede [14] investigated the case of purely vectorial k . Fundamental solutions for the Dirac operator $D + a$, where a is a paravector, are described in [2]. Factorizations of the Helmholtz equation and the related Dirac operators $D + \alpha$, where α is a complex quaternion are investigated by V. V. Kravchenko and M. V. Shapiro [20], [21]. In the context of Maxwell's equations operators generalized Dirac operators considered in [24] by A. McIntosh and M. Mitrea.

After this introduction we give a short overview of the Clifford operator calculus in general, starting with Clifford algebras and then describing Clifford analysis and operator calculus as they needed for applications to equations of mathematical physics independently of a specific Clifford algebra.

In the next section examples are presented, where the main emphasize is laid on the Dirac operator and associated monogenic functions. The last example are parabolic Dirac operators which allows to factorize the heat equation.

Section 3 is concerned with the non-stationary Schrödinger operator and appropriate factorizations by parabolic Dirac operators.

In Section 4 we investigate the factorization of the stationary and non-stationary Schrödinger equation which will lead to the Miura transform.

With the Miura transform as a nonlinear Dirac equation and a procedure based on fix-point principles to solve this nonlinear equation is section 5 concerned.

The last section deals with the Gross-Pitaevskii equation and it's relation to a nonlinear Schrödinger equation and the Miura transform.

2. CLIFFORD OPERATOR CALCULUS

The connection of analysis and algebra has created an deeper understanding of physics and natural sciences. On way to algebra to create analysis can be done by the use of Clifford algebras.

2.1. Clifford algebras. Let $e_1, \dots, e_n$ be an orthonormal basis of $\mathbb{R}^n$. The Clifford algebra $\mathcal{C}_{n,0}$ is the free algebra generated modulo

$$x^2 = -|x|^2 e_0,$$

where e_0 is the identity of the Clifford algebra and the basis elements fulfil the anti-commuting relations:

$$e_i e_j + e_j e_i = \pm 2\delta_{ij} e_0,$$

where δ_{ij} denotes the Kronecker symbol. Let $N = \{1, 2, \dots, n\}$ and for each non-empty subset A of N set

$$e_A = e_{A_1} \dots e_{A_k}, \quad A = \{A_1, \dots, A_k\}, \quad 1 \leq A_1 < A_2 < \dots < A_k \leq n.$$

By convention $e_\emptyset = e_0$ the identity of the Clifford algebra.

There are three operations on a Clifford algebra. The *principal automorphism*

$$e'_A = (-1)^{|A|} e_A, \quad A \subseteq N,$$

where $|A|$ is the cardinality of A , the *principal anti-automorphism*

$$e_A^* = (-1)^{\frac{1}{2}|A|(|A|-1)} e_A,$$

while the *conjugation* is the decomposition of the principal automorphism and anti-automorphism

$$\bar{e}_A = (e_A^*)' = (e_A')^* = (-1)^{\frac{1}{2}|A|(|A|+1)} e_A.$$

It is easily seen that

$$\bar{e}_i = -e_i, \quad i = 1, \dots, n, \quad \text{and} \quad \overline{\bar{e}_i e_j} = \bar{e}_j \bar{e}_i.$$

We will identify the Euclidean space $\mathbb{R}^n$ with $\bigwedge^1 \mathcal{C}_{0,n}$ the space of all vectors of $\mathcal{C}_{0,n}$, i.e. any element $x \in \mathbb{R}^n$ can be identified with

$$x = \sum_{i=1}^n x_i e_i.$$

In this way it is easily seen that each non-zero vector has inverse given by $\frac{-x}{|x|^n}$. More on Clifford algebras and Clifford analysis can be found in the pioneering book [4], the treatment of elliptic boundary value problems is considered in [12]. The function theoretical basis and the application to more complicated partial differential equations like the Navier-Stokes equation is contained in [13]. A (comprehensive) function theory for the Dirac operator is the book [5]. A good description of Clifford analysis and its connection to classical as well global analysis can be found in [9].

2.1.1. Special cases.

1. Complex numbers $\mathbb{C} \sim \mathcal{C}_{0,1}$: Identifying 1 with e_0 and the complex unit i with e_1 . We obtain that the complex numbers $\mathbb{C}$ are equivalent to the Clifford algebra $\mathcal{C}_{0,1}$.

2. Quaternions $\mathbb{H} \sim \mathcal{C}_{0,2}$: Let us denote the basic elements of the quaternions by 1, i , j , k . Then

$$ij = k = -ji, \quad jk = i = -kj, \quad ki = j = -ik,$$

and we identify $1 = e_0$, $i = e_1$, $j = e_2$ and hence $k = e_1 e_2$. Thus the quaternions $\mathbb{H}$ can be identified with $\mathcal{C}_{0,2}$.

Remark 1. Generally Clifford algebras $\mathcal{C}_{p,q}$ are defined by the generating elements e_i , $i = 1, \dots, e_n$, $n = p + q$, where

$$e_i^2 = -1, \quad i = 1, \dots, p, \quad e_i^2 = +1, \quad i = p + 1, \dots, p + q = n, \\ \text{and } e_i e_j + e_j e_i = -2\delta_{ij} e_0.$$

But we restrict ourself to the cases $\mathcal{C}_{0,n}$ and $\mathcal{C}_{n,0}$ because it is possible to factorize all typical types of second order differential operators by using these Clifford algebras.

2.2. Function spaces.

Let $G \subset \mathbb{R}^n$ be a bounded domain with sufficient smooth boundary $\Gamma = \partial G$, whose complement contains a non-empty open set. Then, functions f defined in G with values in $\mathcal{C}_{0,n}$ are considered. These functions may be written as

$$f(x) = \sum_A f_A e_A, \quad x \in G,$$

where $e_A = e_{h_1} \cdots e_{h_k} = e_{h_1 \dots h_k}$, $1 \leq h_1 < \dots < h_k \leq n$, and $e_\emptyset = e_0$. Properties such as continuity, differentiability, integrability and so on, which are ascribed to f have to be possessed by all components $f_A(x)$. In this way the usual Banach spaces of these functions are denoted by C^α , L_p , W_p^k . The norm in the Banach-spaces B is given by

$$\|f\|_B = \sqrt{\sum_A \|f_A\|_B^2}.$$

In particular, we introduce in $L^2(G)$ of *real-valued* function f_A the $\mathcal{C}_{0,n}$ -valued inner product

$$\langle f, g \rangle := \int_G \overline{f(x)} \cdot g(x) dx,$$

where $\overline{f(x)} = \sum_A f_A \bar{e}_A$. The inner product is related to the L^2 -norm by

$$\|f\|_{L^2}^2 = \text{sc} \langle f, f \rangle = \sum_A |f_A|^2(x),$$

where $\text{sc} g = g_0 e_0$ denotes the scalar-part of $g = \sum_A g_A e_A$.

2.3. The operators D , T , F .

In this section we describe in a general way the basic operators which will become specified by the use of a specific Clifford algebra. In general, let G be a domain with smooth boundary $\partial G = \Gamma$ and denote by $n(y)$ the outward pointed normal at $y \in \Gamma$. There are two ways of consideration, the classical one which assumes the domain G to have a Ljapunov boundary Γ , i.e. Γ is of class $C^{1,\alpha}$ with arbitrary $\alpha > 0$ [12], [13]. But these strong assumption on the boundary can be weakened to the consideration of strongly Lipschitz domains [22], [23].

We consider a differential operator D which is a "square root" of a second order differential operator and therefore possesses a fundamental solution $e(x)$, i.e.

$$De(x) = \delta(x) \quad \text{in } G.$$

Because the multiplication in Clifford algebras is not commutative we have to distinguish between left-monogenic functions, i.e. functions f such that $Df = 0$ and right-monogenic functions g , which fulfil $gD = 0$. Then the convolution integral operator

$$Tu = -e \star u = - \int_G e(x-y)u(y) dy$$

is a right but not a left inverse of the operator D and there exists an operator

$$(Fu)(x) = \int_{\Gamma} e(x-y)n(y)u(y) d\Gamma_y, \quad x \in G,$$

such that

$$DT = I \quad \text{and} \quad TD + F = I.$$

The operator F appears by applying the formula of partial integration. By taking limits we obtain

$$\lim_{\text{inside } G / \text{outside } G \rightarrow \Gamma} Fg = \pm g + Sg,$$

with a Hilbert-type operator

$$(Su)(x) = 2 \int_{\Gamma} e(x-y)n(y)u(y) d\Gamma_y, \quad x \in \Gamma.$$

2.4. Decompositions. Another general principal is that

$$L^2(G) = \ker D \oplus \overline{\text{im } D^*}. \quad (1)$$

It is a challenge to prove that the (orthogonal) complement is $\text{im } D^*$, this set is closed itself. Now, it is easy to see that this decomposition defines (orthogonal) projections $\mathbb{P}$ and $\mathbb{Q}$.

Remark 2. *The just mentioned decomposition are orthogonal in L^2 and can be extended to direct sums in L^p , $1 < p < \infty$, and in Sobolev spaces W_p^k , $p \geq 1$.*

The connection between the operator F and the Hilbert-type operator S give raise to another decomposition which is based on the (orthogonal) projections $P = \frac{1}{2}(I + S)$ and $Q = \frac{1}{2}(I - S)$. This is a Hardy-type decomposition

$$L^2(\Gamma) = \text{im } P \oplus \text{im } Q. \quad (2)$$

An interesting connection is given between (1) and (2) in the way that $f \in \ker D$ if and only if $\text{tr } f \in \text{im } P$. This connection is given by the Plemelj-Sokhitzkij's formulas which tell us that $\text{tr } f \in \text{im } P$ if and only $\text{tr } f$ has a monogenic continuation f into G , i.e. $Df = 0$.

2.5. Examples.

2.5.1. Dirac operator. Here, D denotes the Dirac operator

$$Df = \sum_{j=1}^n e_j \frac{\partial f}{\partial x_j},$$

which factorizes the Laplacian:

$$DD = -\Delta_n,$$

where Δ_n denotes the Laplacian in $\mathbb{R}^n$, and we obtain the fundamental solution of the Dirac operator by applying the Dirac operator to the fundamental solution of the Laplacian H . We get the generalized Cauchy kernel

$$e(x) = \frac{-1}{\omega_n} \frac{x}{|x|^n},$$

where ω_n denotes the surface area of the unit sphere in $\mathbb{R}^n$. The first conclusion is

Corollary 1 (Borel-Pompeiu formula). *Let G be a bounded strongly Lipschitz domain of $\mathbb{R}^n$. If $f \in W_p^1(G)$, $1 < p < \infty$, we have*

$$Ff + TDf = \begin{cases} f, & \text{in } G, \\ 0, & \text{in } \mathbb{R}^n \setminus \bar{G}. \end{cases}$$

Similar to the case of one-dimensional complex variables we can define Hardy spaces [25]. Let us denote by $G_+ = G$ and $G_- = \mathbb{R}^n \setminus \bar{G}$.

Definition 1. *For a function $F : G_{\pm} \rightarrow \mathcal{C}_{0,n}$ and $1 < p < \infty$ we set*

$$\|F\|_{\mathcal{H}^p} := \sup_{\delta > 0} \left\{ \int_{\Gamma} |F(x \pm \delta)|^p d\Gamma_x \right\}^{\frac{1}{p}},$$

and define the Hardy spaces of monogenic functions

$$\mathcal{H}^p(G_{\pm}) := \{f \text{ is left monogenic in } G_{\pm}; \|f\|_{\mathcal{H}^p} < \infty\}.$$

There is also a connection to the boundary values of functions from the Hardy spaces $\mathcal{H}^p(G_{\pm})$.

Corollary 2. *Let $1 < p < \infty$. For a left monogenic function H in G , the following are equivalent*

- (1) $H \in \mathcal{H}^p(G)$;
- (2) *There exists $h \in L^p(\Gamma)$ such that H is the (right) Cauchy integral extension of h , i.e. $H(x) = Fh(x)$ for x in G ;*
- (3) *H has a non-tangential boundary limit $H^+(x)$ at almost any point $x \in \Gamma$, i.e. there exists*

$$H^+(x) := \lim_{\{y \in x + \Sigma_{\alpha}, y \rightarrow x\}} H(y) \text{ for a.e. } x \in \Gamma,$$

and H is the (right) Cauchy integral extension of its boundary trace. In addition the norms $\|\cdot\|_{\mathcal{H}^p}$ and $\|H^+\|_{L^p}$ are equivalent, i.e.

$$\|H\|_{\mathcal{H}^p} = \|H^+\|_{L^p}.$$

The most important singular convolution operator is the singular Cauchy integral operator S on Γ , defined for almost all $x \in \Gamma$ by

$$Sf(x) = 2 \lim_{\varepsilon \rightarrow 0} \int_{\{y \in \Gamma: |x-y| > \varepsilon\}} e(x-y)n(y)f(y) d\Gamma_y.$$

Corollary 3 (Plemelj-Sokhotzki's formula). *Let G be a bounded strongly Lipschitz domain of $\mathbb{R}^n$ with boundary $\partial G = \Gamma$. If $f \in L^p(\Gamma)$, $1 < p < \infty$, then*

$$\begin{aligned} (Pf)(x) &= \lim_{\varepsilon \rightarrow 0} \int_{y \in \Gamma: |x-y| > \varepsilon} e(x-y)n(y)f(y) d\Gamma_y + \frac{1}{2}f(x) \\ &= \frac{1}{2}(I + S)f(x). \end{aligned}$$

and hence

$$(Qf)(x) = \frac{1}{2}(I - S)f(x).$$

Moreover, we have

$$Sf = 2Pf(x) - f(x) \quad \text{and} \quad Sf = -2Qf + f.$$

The equality of the integrals is shown in [22]. It follows from Calderón-Zygmund theory. We have

Corollary 4. *The space $L^p(\Gamma)$, $1 < p < \infty$, posses the decomposition*

$$L^p(\Gamma) = \text{im } P \cap L^p(\Gamma) \oplus \text{im } Q \cap L^p(\Gamma),$$

where the decomposition is orthogonal in case $p = 2$ and direct in case $p \neq 2$. Moreover, the spaces $\text{im } P \cap L^p(\Gamma)$ and $\text{im } Q \cap L^p(\Gamma)$ can be identified with the boundary values of the Hardy spaces $\mathcal{H}^p(G_{\pm})$.

The Teodorescu transform is given by

$$Tf(x) = \frac{1}{\omega_n} \int_G \frac{x-y}{|x-y|^n} f(y) dy$$

and from Borel-Pompeiu's formula we immediately conclude that

$$\text{tr } Tf \in \text{im } Q.$$

2.5.2. Cauchy-Riemann operator. Now, D denotes the Cauchy-Riemann operator

$$Df = \frac{\partial f}{\partial x_0} + \sum_{j=1}^n e_j \frac{\partial f}{\partial x_j} \quad \text{and} \quad \bar{D}f = \frac{\partial f}{\partial x_0} - \sum_{j=1}^n e_j \frac{\partial f}{\partial x_j}.$$

Also the Cauchy-Riemann operator factorizes the Laplacian. We have

$$D\bar{D} = \Delta_{n+1},$$

where Δ_{n+1} denotes the Laplacian in $\mathbb{R}^{n+1}$, and we obtain the fundamental solution of the generalized Cauchy kernel

$$e(x) = \frac{1}{\omega_{n+1}} \frac{\bar{x}}{|x|^{n+1}},$$

where ω_{n+1} denotes the surface area of the unit sphere in $\mathbb{R}^{n+1}$.

The Cauchy-Riemann operator can be transform into a Dirac operator by multiplication with e_{n+1} :

$$\frac{\partial f}{\partial x_0} - \sum_{j=1}^n e_j \frac{\partial f}{\partial x_j} e_{n+1} = \frac{\partial f}{\partial x_0} e_{n+1} - \sum_{j=1}^n e_j e_{n+1} \frac{\partial f}{\partial x_j} = \frac{\partial f}{\partial x_0} E_{n+1} - \sum_{j=1}^n E_j \frac{\partial f}{\partial x_j}.$$

On the other hand the Dirac operator can be transformed into the Cauchy-Riemann operator:

$$\sum_{j=1}^n e_j \frac{\partial f}{\partial x_j} (-e_n) = \frac{\partial f}{\partial x_n} + \sum_{j=1}^{n-1} (-e_j e_n) \frac{\partial f}{\partial x_j} = \frac{\partial f}{\partial x_n} + \sum_{j=1}^{n-1} E_j \frac{\partial f}{\partial x_j}.$$

2.5.3. *$\bar{\partial}$ -operator.* The simplest case occurs when the underlying Clifford algebra is just equivalent to the complex numbers, i.e. $\mathcal{C}_{0,1} \sim \mathbb{C}$. In this case we have the complex Cauchy-Riemann operators

$$\bar{\partial}_z = \frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \quad \text{and} \quad \partial_z = \frac{\partial}{\partial x} - i \frac{\partial}{\partial y}$$

which can be identified with

$$D = \frac{\partial}{\partial x} + e_1 \frac{\partial}{\partial y} \quad \text{and} \quad \bar{D} = \frac{\partial}{\partial x} - e_1 \frac{\partial}{\partial y}$$

where $\bar{\partial}$ is identified with D and we have the properties

$$\bar{\partial} \partial = \partial \bar{\partial} = D \bar{D} = \bar{D} D = \Delta = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}.$$

The fundamental solution of the $\bar{\partial}$ -operator is the Cauchy kernel

$$e(z) = -\frac{1}{2\pi i} \frac{1}{z} = \frac{1}{2\pi} \frac{1}{z} i \sim \frac{1}{2\pi} \frac{\bar{z}}{|z|^2} e_1$$

and we have the generalized Cauchy formula (Cauchy-Green formula, Borel formula, ...) as an analog of the Borel-Pompeiu formula:

$$f(z) = \frac{1}{2\pi i} \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta - \frac{1}{\pi} \int_G \frac{\bar{\partial} f(\zeta)}{\zeta - z}, d\zeta.$$

2.5.4. *Generalized Dirac and Cauchy-Riemann operators.* We start with the Dirac operator and consider the disturbed operator

$$D + \alpha = \sum_{j=1}^n e_j \frac{\partial f}{\partial x_j} + \alpha$$

and

$$(D + \alpha)(D - \alpha) = DD - \alpha^2 = -\Delta - \alpha^2,$$

where α is a scalar-valued function. But we can do even more

$$(D + a)(D + \bar{a}) = DD + a\bar{a} = -\Delta - |a|^2.$$

An extensive study of these generalized or disturbed Dirac operators can be found in [20] and [21].

2.5.5. *Parabolic Dirac operators.* Recently (cf. [8]), it was also possible to factorize the heat equation by a Witt-type basis. Which means to add two additional generating elements $\mathfrak{f}$ and $\mathfrak{f}^+$, where

$$\begin{aligned}\mathfrak{f}\mathfrak{f}^+ + \mathfrak{f}^+\mathfrak{f} &= 1, \\ \mathfrak{f}^2 = (\mathfrak{f}^+)^2 &= 0, \\ \mathfrak{f}e_j + e_j\mathfrak{f} &= 0, \\ \mathfrak{f}^+e_j + e_j\mathfrak{f}^+ &= 0.\end{aligned}$$

and

$$\mathbf{D}_{x,t}^\pm = \sum_{j=1}^n e_j \frac{\partial f}{\partial x_j} + \mathfrak{f} \partial_t \pm \mathfrak{f}^+,$$

where

$$(\mathbf{D}_{x,t}^\pm)^2 = -\Delta \pm \partial_t,$$

i.e. $\mathbf{D}_{x,t}^+$ factorizes the heat operator. The fundamental solution for the heat operator is

$$e(x, t) = \frac{H(t)}{(2\sqrt{\pi t})^n} e^{-\frac{|x|^2}{4t}},$$

where $H(t)$ denotes the Heaviside-function. More on parabolic Dirac operators and the application to the non-stationary Navier-Stokes equation can be found in the already mentioned paper [8].

3. PARABOLIC DIRAC OPERATORS RELATED TO THE SCHRÖDINGER EQUATION

Let us denote with C the space-time domain $C = G_t \times [0, \infty) \subset \{(x, t) \in \mathbb{R}^n \times \mathbb{R}^+\}$ and by $\partial C = G \cup M$, where $M = \Gamma_t \times [0, \infty)$, its sufficiently smooth boundary. With the subindex t we emphasis the fact that the space domain (and therefore M) can depend on the time level t , but has to be bounded for all times $t \in [0, \infty)$.

The consideration of the Schrödinger equation is similar to the heat equation but the Schrödinger equations reads as

$$-i\partial_t - \Delta_n,$$

where Δ_n again denotes the Laplacian in $\mathbb{R}^n$. It is easily seen that we obtain a fundamental solution of the Schrödinger equation from the fundamental solution of the heat equation by the transform $t \rightarrow it$, hence

$$E(x, t) = \frac{H(t)}{(2\sqrt{\pi i t})^n} e^{i\frac{|x|^2}{4t}},$$

where $H(t)$ denotes the Heaviside-function.

We consider the operators

$$D_{x,t}^{\pm} = \sum_{j=1}^n e_j \frac{\partial f}{\partial x_j} + \mathfrak{f} \partial_t \pm i \mathfrak{f}^+,$$

and hence

$$\begin{aligned} (D_{x,t}^{\pm})^2 u &= D_{x,t}^{\pm} D_{x,t}^{\pm} u = (D_x + \mathfrak{f} \partial_t \pm i \mathfrak{f}^+) (D_x u + \mathfrak{f} \partial_t u \pm i \mathfrak{f}^+ u) \\ &= D_x D_x u - \mathfrak{f} D_x \partial_t u \mp i \mathfrak{f}^+ D_x u + \mathfrak{f} D_x \partial_t u \\ &\quad + \mathfrak{f}^2 \partial_t^2 u \pm \mathfrak{f} \mathfrak{f}^+ \partial_t u \pm i \mathfrak{f}^+ D_x u + \pm i \mathfrak{f}^+ \mathfrak{f} \partial_t u - (\mathfrak{f}^+)^2 u \\ &= -\Delta u \pm i(\mathfrak{f} \mathfrak{f}^+ + i \mathfrak{f}^+ \mathfrak{f}) \partial_t u = -\Delta u \pm i \partial_t u. \end{aligned}$$

Next, we consider the reduced differential operator

$$D_{x,t} = \sum_{j=1}^n e_j \frac{\partial f}{\partial x_j} + \mathfrak{f} \partial_t.$$

If we introduce the sigma-form

$$d\sigma_{x,t} = D_{x,t} \rfloor dV_x dt$$

we write Stokes theorem as follows

$$\int_{\partial C} f d\sigma_{x,t} g = \int_C (f D_{x,t}) g + f (D_{x,t} g) dV_x dt$$

and thus

$$\int_{\partial C} f d\sigma_{x,t} g = \int_C (f D_{x,t}^+) g + f (D_{x,t}^- g) dV_x dt.$$

From the fundamental solution of the Schrödinger equation we can switch to the fundamental solution of $-\Delta + i \partial_t$ by making a reflection $t \rightarrow -t$ and applying $D_{x,t}^+$ from the right we obtain the fundamental solution

$$\begin{aligned} e(x, t) &= E(x, -t) D_{x,t}^+ = \frac{H(-t)}{(2\sqrt{\pi} i (-t))^n} e^{-i \frac{|x|^2}{4t}} (D_x + \mathfrak{f} \partial_t + i \mathfrak{f}^+) \\ &= \frac{H(-t)}{(2\sqrt{\pi} i (-t))^n} e^{-i \frac{|x|^2}{4t}} \left(\frac{-i}{2t} \sum_{j=1}^n x_j e_j + \mathfrak{f} \left(-\frac{n}{2t} + \frac{i|x|^2}{4t^2} \right) + i \mathfrak{f}^+ \right). \end{aligned}$$

which gives the *Borel-Pompeiu formula* for the operator $D_{x,t}^-$:

$$\int_{\partial C} e(x - x_0, t - t_0) d\sigma_{x,t} g(x, t) = g(x_0, t_0) + \int_C e(x - x_0, t - t_0) (D_{x,t}^- g) dV_x dt$$

and for $g \in \ker D_{x,t}^-$ Cauchy's integral formula:

$$\int_{\partial C} e(x - x_0, t - t_0) d\sigma_{x,t} g(x, t) = g(x_0, t_0).$$

4. FACTORIZATION OF THE SCHRÖDINGER EQUATION

The factorization of the Schrödinger equation is treated in several papers. Mostly the factorization is related to systems of differential equations in mathematical physics [1], [17], [18]. In [19] especially the relationship to the Riccati equation is investigated and quaternionic generalizations of the Riccati equation had are established.

A detailed factorization of the stationary Schrödinger equation for $n = 3, 4$ in an quaternionic context was treated in [3].

The Miura transform can be obtained by factorizing the 1D Schrödinger equation:

$$-\frac{d^2}{dx^2}u - v(x)u = -\left(\frac{d}{dx} + a(x)\right)\left(\frac{d}{dx} - a(x)\right)u.$$

We have

$$\begin{aligned} & -\left(\frac{d}{dx} + a(x)\right)\left(\frac{d}{dx} - a(x)\right)u \\ &= -\left(\frac{d^2u}{dx^2} - \frac{d}{dx}(a(x)u) + a(x)\frac{du}{dx} - a^2(x)u\right) \\ &= -\left(\frac{d^2u}{dx^2} + \left(-\frac{da}{dx} - a^2(x)\right)u\right) \end{aligned}$$

and thus

$$\frac{da}{dx} + a^2(x) = -v(x).$$

4.1. Factorization of the stationary Schrödinger equation. The Helmholtz equation can be treated quite similar to the Laplace equation and can be used to describe Maxwell's equations. The relationship between Maxwell's equations as a system of first order differential equations and the formulation in terms of the Helmholtz equation is nicely characterized by the factorization of the Helmholtz equation. The Helmholtz equation and the factorization in generalized Dirac operators is discussed in [20], [21], [15] and [26]. In higher dimensions it is quite clear that the Helmholtz operator may be factorized by using Dirac operators:

$$-\Delta - k^2 = (D + k)(D - k).$$

More difficult is the case of a variable potential $V_0(x)$. We consider the case $u = u(x)e_0$ and look for suitable functions a with

$$\begin{aligned} (-\Delta - V_0(x))u &= (D + a(x))(D - a(x))u \\ &= DDu - D(a(x)u) + a(x)Du - a^2(x)u \\ &= -\Delta u - \underline{Du \cdot a(x) + a(x)Du} - Da(x) \cdot u - a^2(x)u. \end{aligned}$$

The underlined part does not vanish, because of the non-commutativity of the multiplications of elements of a Clifford algebra. Thus, we will change our approach using a multiplication operator $M^{a(x)}$ defined by

$$M^{a(x)}u(x) := u(x) \cdot a(x).$$

Therefore

$$\begin{aligned} (-\Delta - V_0(x))u &= (D + M^{a(x)}) (D - M^{a(x)}) u \\ &= DDu - D(ua(x)) + Du \cdot a(x) - uDa(x) - ua^2(x) \\ &= -\Delta u - (Da(x) + a^2(x)) u \end{aligned}$$

or

$$Da(x) + a^2(x) = V_0(x). \quad (3)$$

Equation (3) will be called *Miura transform*. It is nonlinear Clifford-valued first-order partial differential equation or equivalently a nonlinear real-valued first-order *system* of partial differential equations.

4.2. Factorization of the non-stationary Schrödinger equation. Now, by using the operator $D_{x,t}^-$ we can also obtain a factorization of the *non-stationary* Schrödinger equation. Let us start with

$$\begin{aligned} (D_{x,t}^- + M^{a(x,t)}) (D_{x,t}^- - M^{a(x,t)}) u(x, t) \\ = D_{x,t}^- D_{x,t}^- u(x, t) - D_{x,t}^- (u(x, t)a(x, t)) + (D_{x,t}^- u(x, t))a(x, t) - u(x, t)a^2(x, t). \end{aligned}$$

We have to evaluate

$$\begin{aligned} D_{x,t}^- (u(x, t)a(x, t)) &= (D_x + \mathfrak{f}\partial_t - i\mathfrak{f}^+) (u(x, t)a(x, t)) \\ &= D_x(u(x, t)a(x, t)) + \mathfrak{f}(\partial_t(u(x, t)a(x, t))) - i\mathfrak{f}^+(u(x, t)a(x, t)) \end{aligned}$$

If we again assume that $u = u(x, t) = u(x, t)e_0$ is a scalar-valued function we can conclude

$$\begin{aligned} &= (D_x u(x, t))a(x, t) + u(x, t)(D_x a(x, t)) + \mathfrak{f}(\partial_t u)a(x, t) + u(x, t)\mathfrak{f}(\partial_t a(x, t)) \\ &\quad - i\mathfrak{f}^+(u(x, t)a(x, t)) \end{aligned}$$

and finally

$$\begin{aligned} (D_{x,t}^- + M^{a(x,t)}) (D_{x,t}^- - M^{a(x,t)}) u(x, t) &= \\ &= D_{x,t}^- D_{x,t}^- u(x, t) - (D_x u(x, t))a(x, t) - u(x, t)(D_x a(x, t)) - \mathfrak{f}(\partial_t u)a(x, t) \\ &\quad - u(x, t)\mathfrak{f}(\partial_t a(x, t)) + i\mathfrak{f}^+(u(x, t)a(x, t)) + (D_{x,t}^- u(x, t))a(x, t) - u(x, t)a^2(x, t) \\ &= (-i\partial_t - \Delta_n)u(x, t) - u(x, t) (D_x a(x, t) + \mathfrak{f}(\partial_t a(x, t)) + a^2(x, t)) \end{aligned}$$

If we compare the last relation with the non-stationary Schrödinger equation we obtain

$$D_x a(x, t) + f(\partial_t a(x, t)) + a^2(x, t) = V(x, t) \quad (4)$$

This equation is similar to the equation

$$\partial_t g + Dg + |g|^2 = 0 \quad (5)$$

which has formal similarities to the canonical Riccati equation. Equation (5) was investigated in [16]. We specify the equation (4) in the following way. In applications the potential usually depends only on x and not on t . Therefore, if we consider a potential $V = V(x)$ equation (4) suggest to consider also functions $a = a(x)$ which depend only on x . Which implies that we will consider vector-valued functions $\underline{a}(x)$ depending only on x and we end up with the equation

$$D_x \underline{a}(x) + \underline{a}^2(x) = V(x), \quad (6)$$

which is the same as for the factorization of the stationary Schrödinger equation.

5. THE NONLINEAR DIRAC EQUATION

In this section we want to solve equation (6) under the following assumptions:

- (1) the potential $V = V(x)$ is a scalar-valued function that depends only on $x \in G$, where G is a compact and smooth domain of $\mathbb{R}^n$.
- (2) we are looking for a vector-valued function $\underline{a}(x) = \sum_{i=1}^n a_i(x) e_i$ that depends only on $x \in G$.

If we apply the T -operator and use Borel-Pompeiu's formula (cf. Corollary 1) we obtain

$$\underline{a} - F\underline{a} = T(V + |\underline{a}|^2),$$

because we are only looking for $\underline{a} \in \text{im } Q$ we have $F\underline{a} = 0$ and we end up with

$$\underline{a} = T(V + |\underline{a}|^2)$$

Our iteration procedure reads now as follows:

$$\underline{a}_n := T(V + |\underline{a}_{n-1}|^2), \quad n = 1, 2, \dots,$$

Obviously, $V(x) + |\underline{a}_{n-1}|^2 \in \mathbb{C}$ reproduces $\underline{a}_n = T(V + |\underline{a}_{n-1}|^2) \in \bigwedge^1 \mathcal{C}_{0,n}$ and because of $\text{tr } T \in \text{im } Q$, also $\text{tr } \underline{a}_n \in \text{im } Q$.

That means that we also have to assume $\text{tr } \underline{a}_{n-1} \in \text{im } Q$.

Now, we consider the regularity. The Teodorescu transform is a weakly singular integral operator and therefore

$$V \in L^p(G) \Rightarrow TV \in W_p^1(G), \quad 1 < p < \infty,$$

on the other hand Sobolev's embedding theorems $W_p^1(G) \hookrightarrow L^{p^*}(G)$, where $p^* \leq \frac{np}{n-p} \leq n$, with embedding constant C leads to the following. Set $p^* = 2p$ and assume that $\underline{a}_{n-1} \in W_p^1(G)$. Then

$$\begin{aligned}
\| |\underline{a}_{n-1}|^2 \|_{L^p} &= \left(\int_G |\underline{a}_{n-1}(x)|^{2p} dG \right)^{1/p} = \left(\int_G \left(\sum_{j=1}^n a_{j,n-1}^2(x) \right)^{\frac{1}{2} \cdot 2p} dG \right)^{1/p} \\
&= \left(\int_G \left(\sum_{j=1}^n a_{j,n-1}^2(x) \right)^p dG \right)^{1/p} = \left\| \sum_{j=1}^n a_{j,n-1}^2(x) \right\|_{L^p} \\
&\leq \sum_{j=1}^n \| a_{j,n-1}^2(x) \|_{L^p} = \sum_{j=1}^n \left(\int_G a_{j,n-1}^{2p}(x) dG \right)^{1/p} \\
&= \sum_{j=1}^n \| a_{j,n-1}(x) \|_{L^{2p}}^2 \leq \sum_{j=1}^n C^2 \| a_{j,n-1}(x) \|_{W_p^1}^2 = C^2 \| \underline{a}_{n-1}(x) \|_{W_p^1}^2,
\end{aligned}$$

i.e. $\underline{a}_{n-1} \in W_p^1(G)$ implies $|\underline{a}_{n-1}|^2 \in L_p(G)$ and finally also $T|\underline{a}_{n-1}|^2 \in W_p^1(G)$.

Thus we assume that $n > p$ and

$$\frac{np}{n-p} \geq 2p \iff np \geq 2np - p^2 \iff 2p^2 \geq np \iff 2p \geq n \iff p \geq \frac{n}{2}.$$

Therefore we need the following assumptions too:

- (1) $n > p \geq \frac{n}{2} > 1$, if $n > 2$ (or $n > p > 1$ if $n = 2$), $p \in \mathbb{R}$,
- (2) $\underline{a}_{n-1} \in W_p^1(G)$ and $\text{tr } \underline{a}_{n-1} \in \text{im } Q \cap W_p^{1-\frac{1}{p}}(\Gamma)$.

The last assumption involves that if $\underline{a}_{n-1}$ belongs to $W_p^1(G)$ due to the trace theorem for Sobolev spaces $\text{tr } \underline{a}_{n-1}$ belongs to $W_p^{1-\frac{1}{p}}(\Gamma)$.

In order to apply fixed-point theorems we first prove the boundedness of the sequence $\{\underline{a}_n\}$. We have

$$\begin{aligned}
\| \underline{a}_n \|_{W_p^1} &\leq \| T \|_{[L^p, W_p^1]} (\| V \|_{L^p} + \| |\underline{a}_{n-1}|^2 \|_{L^p}) \\
&\leq k_1 \left(\| V \|_{L^p} + C^2 \| \underline{a}_{n-1} \|_{W_p^1}^2 \right) \\
&\leq k_1 \| V \|_{L^p} + k_2 \| \underline{a}_{n-1} \|_{W_p^1}^2,
\end{aligned} \tag{7}$$

where $k_1 = \| T \|_{[L^p, W_p^1]}$, $k_2 = k_1 C^2$, and C is again the embedding constant for $W_p^1 \hookrightarrow L^{2p}$.

Lemma 1. *We assume that $\| V \|_{L^p} \leq \frac{1}{4k_1 k_2}$ and denote by*

$$W = \sqrt{\frac{1}{4k_2^2} - \frac{k_1}{k_2} \| V \|_{L^p}}. \tag{8}$$

Then

$$\frac{1}{2k_2} - W \leq \| \underline{a}_{n_0} \|_{W_p^1} \leq \frac{1}{2k_2} + W$$

implies that $\|\underline{a}_{n_0+1}\|_{W_p^1} \leq \|\underline{a}_{n_0}\|_{W_p^1}$ and the sequence $\{\underline{a}_n\}_{n \geq n_0}$ will be bounded by $\|\underline{a}_{n_0}\|_{W_p^1}$.

Proof. The inequality (8) can be rewritten as

$$-W \leq \|\underline{a}_{n_0}\|_{W_p^1} - \frac{1}{2k_2} \leq W$$

and hence is equivalent to

$$\|\underline{a}_{n_0}\|_{W_p^1}^2 - \frac{1}{k_2} \|\underline{a}_{n_0}\|_{W_p^1} + \frac{1}{4k_2^2} \leq W^2 = \frac{1}{4k_2^2} - \frac{k_1}{k_2} \|V\|_{L^p}$$

and we obtain that

$$\|\underline{a}_{n_0}\|_{W_p^1}^2 - \frac{1}{k_2} \|\underline{a}_{n_0}\|_{W_p^1} + \frac{k_1}{k_2} \|V\|_{L^p} \leq 0$$

and using (7)

$$\|\underline{a}_{n_0+1}\|_{W_p^1} \leq k_1 \|V\|_{L^p} + k_2 \|\underline{a}_{n_0}\|_{W_p^1}^2 \leq \|\underline{a}_{n_0}\|_{W_p^1}.$$

□

Theorem 1. Suppose that $n > p \geq \frac{n}{2} > 1$, $p \in \mathbb{R}$, and $\|V\|_{L^p} \leq \frac{1}{4k_1k_2}$. Then the equation

$$\underline{a} = T(V + |\underline{a}|^2)$$

has at least one solution with

$$\|\underline{a}\|_{W_p^1} \leq \frac{1}{2k_2} + \sqrt{\frac{1}{4k_2^2} - \frac{k_1}{k_2} \|V\|_{L^p}}.$$

Proof. From 1 we immediately see that

$$\|\underline{a}_{n-1}\|_{W_p^1} \leq \frac{1}{2k_2} + W \Rightarrow \|\underline{a}_n\|_{W_p^1} \leq \frac{1}{2k_2} + W.$$

If we start with $\underline{a}_0 \in W_p^1(G)$ such that $\|\underline{a}_0\|_{W_p^1} \leq \frac{1}{2k_2} + W$ then the sequence $\{\underline{a}_n\}_{n \in \mathbb{N}}$ is bounded from above by $\frac{1}{2k_2} + W$. Hence there exists a subsequence $\{\underline{a}_{n'}\} \subset W_p^1(G)$ with $\underline{a}_{n'} \rightharpoonup \underline{a}$ as $n' \rightarrow \infty$. Due to the continuity of $T : L^p(G) \rightarrow W_p^1(G)$ we conclude

$$\underline{a} = T(V + |\underline{a}|^2).$$

The norm estimates follows from the weak convergence of $\underline{a}_{n'}$ in a convex set. □

But we can do better

Theorem 2. We assume that

- (1) $n > p \geq \frac{n}{2} > 1$, $p \in \mathbb{R}$,
- (2) $\|V\|_{L^p} \leq \frac{1}{4k_1k_2}$,
- (3) $\underline{a}_0 \in W_p^1(G)$ and $\text{tr } \underline{a}_0 \in \text{im } Q \cap W_p^{1-1/p}(\Gamma)$,
- (4) $\|\underline{a}\|_{W_p^1} \leq \frac{1}{2k_2} - \sqrt{\frac{1}{4k_2^2} - \frac{k_1}{k_2} \|V\|_{L^p}}$.

and define the sequence $\{\underline{a}_n\}_{n \in \mathbb{N}}$ by

$$\underline{a}_n = T(V + |\underline{a}_{n-1}|^2), \quad n = 1, 2, \dots$$

Then there exists a unique solution $\underline{a} \in W_p^1(G)$ with $\text{tr } \underline{a} \in \text{im } Q \cap W_p^{1-1/p}(\Gamma)$ of

$$\underline{a} = T(V + |\underline{a}|^2).$$

Moreover, the sequence $\{\underline{a}_n\}_{n \in \mathbb{N}}$ converges to $\underline{a}$ in W_p^1 and the solution $\underline{a}$ fulfils the estimate

$$\|\underline{a}\|_{W_p^1} \leq \frac{1}{2k_2} - \sqrt{\frac{1}{4k_2^2} - \frac{k_1}{k_2} \|V\|_{L^p}}.$$

Proof. To apply Banach's fix-point theorem we investigate the contractivity of the mapping $T(v + |\underline{a}|^2)$. At first we get

$$\|\underline{a}_n - \underline{a}_{n-1}\|_{W_p^1} = \|T(|\underline{a}_{n-1}|^2 - |\underline{a}_{n-2}|^2)\|_{W_p^1} \leq k_1 \| |\underline{a}_{n-1}|^2 - |\underline{a}_{n-2}|^2 \|_{L^p}.$$

Furthermore,

$$\begin{aligned} \| |\underline{a}_{n-1}|^2 - |\underline{a}_{n-2}|^2 \|_{L^p} &= \| (|\underline{a}_{n-1}| - |\underline{a}_{n-2}|)(|\underline{a}_{n-1}| + |\underline{a}_{n-2}|) \|_{L^p} \\ &\leq \| |\underline{a}_{n-1}| - |\underline{a}_{n-2}| \|_{L^{2p}} \| |\underline{a}_{n-1}| + |\underline{a}_{n-2}| \|_{L^{2p}} \\ &\leq \| |\underline{a}_{n-1}| - |\underline{a}_{n-2}| \|_{L^{2p}} \| |\underline{a}_{n-1}|^2 \|_{L^p}^{1/2} + \| |\underline{a}_{n-2}|^2 \|_{L^p}^{1/2} \\ &\leq \| |\underline{a}_{n-1}| - |\underline{a}_{n-2}| \|_{L^{2p}} \left(C \|\underline{a}_{n-1}\|_{W_p^1} + C \|\underline{a}_{n-2}\|_{W_p^1} \right) \\ &\leq 2C \| |\underline{a}_{n-1}| - |\underline{a}_{n-2}| \|_{L^{2p}} \left(\frac{1}{2k_2} - \sqrt{\frac{1}{4k_2^2} - \frac{k_1}{k_2} \|V\|_{L^p}} \right) \end{aligned}$$

Now, we use that $\underline{a}_{n-1}$ and $\underline{a}_{n-2}$ are vector-valued and conclude

$$\| |\underline{a}_{n-1}| - |\underline{a}_{n-2}| \|_{L^{2p}} \leq \| |\underline{a}_{n-1} - \underline{a}_{n-2}| \|_{L^{2p}} \leq \| |\underline{a}_{n-1} - \underline{a}_{n-2}|^2 \|_{L^p}^{1/2} \leq C \| \underline{a}_{n-1} - \underline{a}_{n-2} \|_{W_p^1}.$$

Then the contractivity constant can be bound from above by

$$L = 2k_1 C^2 \left(\frac{1}{2k_2} - W \right) = 2k_2 \left(\frac{1}{2k_2} - W \right) = 1 - 2k_2 W = 1 - \sqrt{1 - 4k_1 k_2 \|V\|_{L^p}}.$$

□

Remark 3. Actually, it is no problem to find a suitable $\underline{a}_0 \in W_p^1(G)$ and $\text{tr } \underline{a}_0 \in \text{im } Q \cap W_p^{1-1/p}(\Gamma)$. Obviously, $\underline{a} = \vec{0}$ fulfils all necessary conditions. But we have also other possibilities. Choose an arbitrary scalar-valued function $b(x)$ such that $\|b\|_{L^p}$ is small enough then $\underline{a}_0 = Tb$ is obviously vector-valued and fulfils all necessary conditions.

Remark 4. More serious problem is that we had assumed the potential to be of small L^p -norm which need not to be fulfilled. In general the equation

$$\underline{a} = T(|\underline{a}|^2) + TV$$

is a system of nonlinear Fredholm equations of second with a weakly singular integral operator T . Hence methods to solve such types of nonlinear Fredholm equations could be applied.

In what follows we will see that the solution of 6 is closely related to the solution of the linear Schrödinger equation.

The following considerations are an easy generalization from the three dimensional case which was treated in [19] to the n -dimensional one. We go back to the Miura transform which could be considered as a Riccati equation as done in [19]. The Riccati equation

$$\partial u = pu^2 + qu + r,$$

where p, q, r are functions, has received a lot of attention especially because of the wide range of applications in which it appears. This equation can be reduced into its canonical form

$$\partial w + w^2 = -v.$$

Also the Riccati equation is related to the one-dimensional Schrödinger equation

$$-\Delta u - vu = 0 \tag{9}$$

with a function v by the substitution

$$w = \frac{\partial u}{u} = \partial \ln u.$$

Therefore the logarithmic derivative (in the sense of Marchenko) of a scalar-valued function u such that $u \neq 0$ in G , is defined as

$$\check{\partial} u = u^{-1} Du = D(\ln u).$$

Proposition 1 ((cf. [19])). *The scalar-valued function φ is a solution of the Schrödinger equation (9) if and only if $\underline{a} := \check{\partial}\varphi$ is a solution of the Miura transform*

$$D\underline{a} = v + |\underline{a}|^2.$$

Proof. Suppose that there exists a function φ such that $\underline{a} = \check{\partial}\varphi$. Then

$$D\underline{a} = \frac{1}{\varphi^2} |D\varphi|^2 - \frac{1}{\varphi} \Delta\varphi, \quad \text{and} \quad |\underline{a}|^2 = \frac{1}{\varphi^2} |D\varphi|^2,$$

and thus

$$D\underline{a} - |\underline{a}|^2 = -\frac{1}{\varphi} \Delta\varphi = v \iff -\Delta\varphi - v\varphi = 0.$$

Conversely, if φ is a solution of the Schrödinger equation $\underline{a} = \check{\partial}\varphi$ is a solution of the Miura transform. $\square$

6. APPLICATION TO THE GROSS-PITAEVSKII EQUATION

Our considerations of the physical background of the Gross-Pitaevskii equation is based on the paper [31] and [32]. The Gross-Pitaevskii equation for the order parameter of the mean field theory is derived by the Fermi's zero-range pseudo-potential [10] and [30].

6.1. Bose-Einstein condensate. An ideal Bose gas is a quantum-mechanical version of a classical ideal gas. It is composed of bosons, which have an integral value of spin, and obey Bose-Einstein statistics. The statistical mechanics of bosons were developed by Satyendra Nath Bose for photons, and extended to massive particles by Albert Einstein who realized that an ideal gas of bosons would form a condensate at a low enough temperature, unlike a classical ideal gas. This condensate is known as a Bose-Einstein condensate.

In dilute gases, only binary collisions are taken into account. The Hamiltonian of the interacting system is:

$$\hat{H} = \int dx \hat{\Psi}^\dagger(x) \left[-\frac{\hbar^2}{2m} \Delta + V(x) \right] \hat{\Psi}(x) + \frac{1}{2} \int dx dx' \hat{\Psi}^\dagger(x) \hat{\Psi}^\dagger(x') U(x-x') \hat{\Psi}(x') \hat{\Psi}(x),$$

where $\hat{\Psi}$ is the annihilation operator, m the mass of a single boson, $\hbar$ reduced Planck constant. Experimental realizations of Boson-Einstein condensate with dilute atomic gases has been achieved in a trapping potential $V(x)$, whose shape is harmonic.

The time evolution of the condensate is given by the Heisenberg equation

$$\begin{aligned} i\hbar \frac{\partial}{\partial t} \hat{\Psi}(x, t) &= [\hat{\Psi}, \hat{H}] \\ &= \left[-\frac{\hbar^2}{2m} \Delta + V(x) + \int \hat{\Psi}^\dagger(x', t) U(x-x') \hat{\Psi}(x', t) \right] \hat{\Psi}(x, t). \end{aligned}$$

The mean-field approximation consists in replacing $\hat{\Psi}$ by the classical field Ψ_0 . In the integral containing $U(x'-x)$, this is a bad approximation for short distances. However, only collisions at low energy are relevant, which can be described by a single parameter: the s-wave scattering length α . Then one can write:

$$U(x'-x) = g\delta(x-x') \quad \text{with} \quad g = \frac{4\pi \hbar^2 \alpha}{m}.$$

And we obtain the *Gross-Pitaevskii equation*

$$i\hbar \frac{\partial}{\partial t} \Psi_0(x, t) = \left(-\frac{\hbar^2}{2m} \Delta + V(x) + g|\Psi_0(x, t)|^2 \right) \Psi_0(x, t),$$

which is valid if

- the number of atoms in the condensate $\gg 1$,
- low temperature
- $\alpha \ll$ average distance between atoms: $\bar{\nu}\alpha^3 \ll 1$ (ν is the density of the gas).

Here, we have the Fermi pseudo-potential (given by the Dirac distribution) has one parameter – the scattering length. Thus, the simplest model which can describe nonlocal repulsive interactions involves introduction of an effective potential with two parameters: the strength of interaction and the interaction range. The symmetry properties and the assumption of short-range interactions is enough to single out the effective potential with two parameters. For spherically symmetric interaction the U depends only on the relative distance $U(x - x') = U(|x - x'|)$. Its Fourier image is then a real function of the squared wave number $\mathbf{k}$:

$$\hat{U} = \hat{U}(\mathbf{k}^2).$$

For the Fermi pseudo-potential is $\hat{U} = 1$. In the next order of approximation we take the *Lorentzian*:

$$\hat{U} = (1 + \alpha^2 \mathbf{k}^2)^{-1}.$$

The Lorentzian has been used recently as an approximation of the finite-range potential. In three dimensions the Lorentzian corresponds to the *Yukawa* effective potential:

$$U_{3D} = \frac{1}{4\pi\alpha^2 x} \exp\left(-\frac{x}{\alpha}\right),$$

and in two dimensions the effective interaction potential is given by

$$U_{2D} = \frac{1}{2\pi\alpha^2} K_0\left(\frac{x}{\alpha}\right),$$

where $K_0(z)$ is the MacDonald's function.

In the following we will use the operator form for the effective potential:

$$\Lambda = 1 - \alpha^2 \nabla^2 = 1 - \alpha^2 \Delta.$$

Which allows to consider the nonlocal Gross-Pitaevskii equation as a system with an additional dependent variable F describing the nonlinear term:

$$(1 - \alpha^2 \nabla^2)F = |\psi|^2, \tag{10}$$

$$\left(i\hbar \frac{\partial}{\partial t} + \frac{\hbar^2}{2m} \Delta - V(x) - gF\right) \Psi_0(x, t) = 0. \tag{11}$$

6.2. Stationary Gross-Pitaevskii equation.

We obtain the *stationary Gross Pitaevskii equation* by setting

$$\Psi_0(x, t) = \varphi(x) e^{-i\mu t/\hbar}, \quad \text{where } \mu \text{ is the chemical potential,}$$

and then

$$\left(-\frac{\hbar^2}{2m}\Delta + V(x) + g|\varphi(x)|^2\right)\varphi(x) = \mu\varphi(x), \quad \text{or} \quad (1 - \alpha^2\nabla^2)F(x) = |\varphi(x)|^2, \quad (12)$$

$$\left(-\frac{\hbar^2}{2m}\Delta + gF(x) + V(x) - \mu\right)\varphi(x) = 0. \quad (13)$$

Where the nonlinear equation is equivalent to a system of linear equations. Here, we see that the stationary Gross-Pitaevskii equation fits perfect in our schema. Equation (12) is a Helmholtz equation

$$\left(-\Delta - \frac{1}{(i\alpha)^2}\right)F(x) = \frac{1}{\alpha^2}|\varphi(x)|^2$$

with constant wave number and equation (13) is a (stationary) Schrödinger equation

$$\left(-\Delta - \frac{-2m}{\hbar^2}(gF(x) + V(x) - \mu)\right)\varphi(x) = 0,$$

of which φ is a solution if and only if

$$\underline{a} = \check{\partial}\varphi = \varphi^{-1}D\varphi = D(\ln \varphi)$$

is a solution of the Miura transform

$$D\underline{a} - |\underline{a}|^2 = \frac{-2m}{\hbar^2}(gF(x) + V(x) - \mu).$$

6.3. Non-stationary Gross-Pitaevskii equation.

Also the non-stationary Gross-Pitaevskii equation

$$i\hbar\frac{\partial}{\partial t}\psi(x, t) = \left(-\frac{\hbar^2}{2m}\Delta + V(x) + g|\psi(x, t)|^2\right)$$

maybe written as a system of linear equations:

$$(1 - \alpha^2\nabla^2)F(x, t) = |\psi(x, t)|^2, \quad (14)$$

$$\left(\left(-i\hbar\frac{\partial}{\partial t} - \frac{\hbar^2}{2m}\Delta\right) + V(x) + gF(x, t)\right)\psi(x, t) = 0. \quad (15)$$

Equation (14) is again equivalent to a Helmholtz equation with constant wave number but the second equation (15) has due to the part $F(x, t)$ a potential which depends on the time t .

REFERENCES

- [1] Bernstein, S., 1996, Factorization of Solutions of the Schrödinger equation. In: Sprößig, W. and Gürlebeck, K. (eds) *Proceedings of the Symposium "Analytical and Numerical Methods in Quaternionic and Clifford Analysis" held at Seiffen, Germany 1996*, University of Mining and Technology Freiberg, 207–216.
- [2] Bernstein, S., 1996, Fundamental solutions for Dirac-type operators. In: J. Ławrynowicz (ed.), *Generalizations of Complex Analysis and their Applications in Physics*, Banach Center Publ. vol. 37, 159–172.
- [3] Bernstein, S. and Gürlebeck, K., 1999, On a higher-dimensional Miura transform. *Complex Variables, Theory Appl.* 38, No.4, 307–319.
- [4] Brackx, F., Delanghe, R. and Sommen, F., 1982, *Clifford analysis*. Research Notes in Mathematics, 76. Boston - London - Melbourne: Pitman Advanced Publishing Program. X.
- [5] Delanghe, R., Sommen, F. and Souček, V., 1992, *Clifford algebra and spinor-valued functions. A function theory for the Dirac operator*. Related REDUCE software by F. Brackx and D. Constaes. Mathematics and its Applications 53, Dordrecht etc.: Kluwer Academic Publishers.
- [6] Brackx, F. and Van Acker, N., 1993, Boundary value theory for eigenfunctions of the Dirac operator. *Bull. Soc. Math. Belg.*, vol. 45, # 2, Ser. B, 113–123.
- [7] Brackx, F., Delanghe, R., Sommen, F. and Van Acker, N., 1993, Reproducing kernels on the unit sphere. In: Pathak, R. S. (ed.), *Generalized functions and their applications*. Proceedings of the international symposium, held December 23–26, 1991 in Varanasi, India. New York, Plenum Press, 1–10.
- [8] Cerejeiras, P., Kähler, U. and Sommen, F., to appear, Parabolic Dirac operators and the Navier-Stokes equations over time-varying domains.
- [9] Gilbert, J. E. and Murray, M. A. M., 1991, *Clifford algebras and Dirac operators in harmonic analysis*. Cambridge studies in advanced mathematics 26, Cambridge university Press, Cambridge.
- [10] Gross, E. P., 1961, *Nuovo Cimento*, 20, 454; 1963, *J. Math. Phys.* 4, 195.
- [11] Gürlebeck, K., 1986, Hypercomplex Factorization of the Helmholtz equation. *Zeitschrift für Analysis und ihre Anwendungen*, Bd. 5(2), 125–131.
- [12] Gürlebeck, K. and Sprößig, W., 1990, *Quaternionic Analysis and Elliptic Boundary Value Problems*. Birkhäuser Verlag, Basel.
- [13] Gürlebeck, K. and Sprößig, W., 1997, *Quaternionic and Clifford calculus for physicists and engineers*. Mathematical Methods in Practice. Chichester: Wiley.
- [14] Huang, L., 1990, The existence and uniqueness theorems of the linear and nonlinear R.-H. problems for the generalized holomorphic vector of the second kind. *Acta Math. Sci. Engl. Ed.* 10, no. 2, 185–199.
- [15] Kravchenko, V. V. and Santana, G., 1996, On singular integral operators associated with the vector Helmholtz equation. *J. Nat. Geom.* 10, No.2, 119–136.
- [16] Kravchenko, V. G. and Kravchenko, V. V., 1994, On some nonlinear equations generated by Fueter type operators, *Zeitschrift für Analysis und ihre Anwendungen*, 13, 599–602.
- [17] Kravchenko, V. G. and Kravchenko, V. V., 2003, Quaternionic factorization of the Schrödinger operator and its applications to some first-order systems of mathematical physics. *J. Phys. A, Math. Gen.* 36, No.44, 11285–11297.
- [18] Kravchenko, V. G. and Kravchenko, V. V., 2004, On a generalized Schrödinger operator and Darboux transformation. In: Simos, T. E. (ed.) et al., *ICNAAM 2004. International conference on numerical analysis and applied mathematics 2004, Chalkis, Greece, September 10-14, 2004. Official conference of the European Society of Computational Methods in Science and Engineering (ESCMSE)*. Weinheim: Wiley-VCH. 502–505, 2004.

- [19] Kravchenko, V. G., Kravchenko, V. V. and Williams, B., 2001, A quaternionic generalization of the Riccati differential equation. In: Brackx, F. (ed.) et al., *Clifford analysis and its applications*. Proceedings of the NATO advanced research workshop, Prague, Czech Republic, October 30–November 3, 2000, Dordrecht: Kluwer Academic Publishers. NATO Sci. Ser. II, Math. Phys. Chem. 25, 143–154.
- [20] Kravchenko, V. V. and Shapiro, M. V., 1994, Helmholtz operator with a quaternionic wave number and associated function theory. *Deformations of Mathematical Structures, II*, Kluwer Academic Publishers (ed.: J. Ławrynowicz), 101–128.
- [21] Kravchenko, V. V. and Shapiro, M. V., 1996, *Integral representations for spatial models of mathematical physics*, Pitman Research Notes in Mathematics Series 351, Pitman.
- [22] Li, C., McIntosh, A. and Semmes, S., 1992, Convolution singular integrals on Lipschitz surfaces. *Journal of the American Mathematical Society* 5, 455–481.
- [23] Li, C., McIntosh, A. and Qian, T., 1994, Clifford algebras, Fourier transforms, and singular convolution operators on Lipschitz surfaces. *Revista Matemática Iberoamericana*, 10, 665–721.
- [24] McIntosh, A. and Mitrea, M., 1999, Clifford algebras and Maxwell’s equations in Lipschitz domains. *Math. Methods Appl. Sci.* 22, No.18, 1599–1620.
- [25] Mitrea, M., 1994, *Clifford Wavelets, Singular Integrals and Hardy Spaces*, Lecture Notes in Mathematics 1575, Springer Verlag Berlin Heidelberg.
- [26] Mitrea, M., 1996, Boundary value problems and Hardy spaces associated to the Helmholtz equation in Lipschitz domains, *J. Math. Anal. Appl.* 202, No. 3, 819–842.
- [27] Miura, R. M., 1968, Korteweg–de Vries equations and generalizations I. A remarkable explicit nonlinear transformation. *J. Math. Phys.*, 9, 1202–1204.
- [28] Obolashvili, E., 1975, Space generalized holomorphic vectors. *Diff. Urav.* T. XI. 1, 108–115.
- [29] Obolashvili, E., 1988, Effective solutions of some boundary value problems in two and three cases. *Functional analytical methods in complex analysis and applications to PDE*, Trieste, 149–172.
- [30] Pitaevskii, L. P., 1961, *Zh. Eksp. Teor. Fis.*, 40, 646 [Sov. Phys. JETP 13 451].
- [31] Shchesnovich, V. S. and Kraenkel, R. A., 2004, Vortices in nonlocal Gross-Pitaevskii equation. *J. Phys. A, Math. Gen.* 37, No.26, 6633–6651.
- [32] Sigwarth, O., 2004, Gross-Pitaevskii equation in atomic Bose-Einstein condensates, <http://www.phy.uni-bayreuth.de/~btpj03/gpeq.pdf>.
- [33] Xu, Z., 1991, A function theory for the operators $D - \lambda$. *Complex Variables Theory and Appl.* 16: 27–42.
- [34] Xu, Z., 1992, Helmholtz equations and boundary value problems. In: *Partial differential equations with complex analysis* (H. Begehr, A. Jeffrey, eds), Pitman Research Notes in Mathematics Series 262, 204–214.

INSTITUTE OF APPLIED ANALYSIS, UNIVERSITY OF MINING AND TECHNOLOGY, 09599 FREIBERG, GERMANY, E-MAIL: SWANHILD.BERNSTEIN@TU-FREIBERG.DE .